



Résumés

ELLIPSOMETRIE FEMTOSECONDE : Aspects techniques et applications

Le 24 mars 2020
Au FIAP, 30 rue Cabanis, 75014 Paris



Lundi 9 décembre 2019

| | |
|---------------|---|
| 09:15 - 09:45 | <i>Accueil / Café</i> |
| 09:45 - 10:30 | Chiralité et Dichroïsme circulaire vibrationnel - Thierry Buffeteau |
| 10:30 - 11:15 | Dichroïsme circulaire de photoélectron - Yann Mairesse |
| 11:15 - 11:30 | <i>Pause café</i> |
| 11:30 - 12:15 | Dichroïsme circulaire électronique femtoseconde - Malte Oppermann |
| 12:15 - 13:30 | <i>Déjeuner</i> |
| 13:30 - 14:15 | Imagerie Polarimétrique - Marie-Claire Schanne-Klein |
| 14:15 - 15:00 | Femtomagnétisme - Nicolas Bergéard |
| 15:00 - 15:15 | <i>Pause café</i> |
| 15:15 - 16:00 | Faisceaux lumineux à moment orbital angulaire – Thierry Ruchon |
| 16:00 - 16:30 | Discussions - Bilan et perspectives utilisateurs |

Dichroïsme Circulaire Vibrationnel (VCD) : Origine, mesure et exemples d'application

Thierry Buffeteau

*Institut des Sciences Moléculaires (UMR 5255 – CNRS), Université de Bordeaux,
33405 Talence, France
e-mail: thierry.buffeteau@u-bordeaux.fr*

L'origine du dichroïsme circulaire vibrationnel (VCD) et ses particularités par rapport à l'absorption infrarouge conventionnelle seront présentées au cours de cet exposé. Après avoir défini les grandeurs pertinentes, nous verrons comment le signal VCD peut être mesuré expérimentalement et comment les spectres VCD peuvent être calculés à partir de calculs de chimie quantique. Quelques exemples d'application du VCD seront ensuite discutés : la détermination de la configuration et de la conformation absolue de molécules chirales, la mise en évidence de la chiralité dans des systèmes moléculaires organisés, l'utilisation du VCD pour identifier et quantifier les structures secondaires de polypeptides ou de protéines en solution ou déposés en films minces ...

Femtosecond and attosecond dynamics probed by photoelectron circular dichroism

Yann Mairesse

*CELIA, Université de Bordeaux,
33405 Talence, France
e-mail: yann.mairesse@celia.u-bordeaux.fr*

When randomly oriented chiral molecules are ionized by circularly polarized light, the photoelectrons are preferentially ejected forward or backward the light propagation axis, depending on the handedness of the molecule and light. This chiroptical process, called photoelectron circular dichroism, is remarkably intense and sensitive. I will show how it can be used to track femtosecond dynamics of chiral molecules, as well as how the attosecond dynamics underlying the chiral photoionization process can be measured.

Capturing the ultrafast evolution of excited state chirality with time-resolved electronic circular dichroism

Malte Oppermann

Ecole Polytechnique Fédérale de Lausanne & LACUS, Lausanne, Switzerland
e-mail: malte.oppermann@epfl.ch

For understanding the functioning of many photo-driven molecular machines, it is crucial to capture the temporal evolution of their chirality and thus structure upon photo-excitation. This is especially relevant for understanding many biological systems, such as photo-active proteins and the kinetics of novel synthetic machines, such as unidirectional molecular motors. Steady-state electronic circular dichroism (CD) has become a standard analytical tool to obtain solution-phase structural information via a molecule's chiral properties. Especially in the deep ultraviolet (UV) spectral range (< 300 nm), it is sensitive to the UV-transitions in proteins and many organic ligands in functional chiral complexes. However, pushing CD spectroscopy into the time-domain has remained a challenge, with only few isolated reports with sub-nanosecond resolution [1].

In this context, I will present a novel broadband time-resolved CD spectrometer in the deep-UV region (250–370 nm) with sub-picosecond time-resolution [2]. With this instrument, it is possible to extract broadband CD spectra of photo-excited states and track their temporal evolution. From the characterisation of the structurally well-defined photo-excited triplet metal-to-ligand charge transfer (MLCT) state in $[\text{Ru}(\text{bpy})_3]^{2+}$, I will proceed to the more complicated dynamics of photo-excited supramolecular systems: $[\text{Fe}(\text{bpy})_3]^{2+}$ is configurationally labile in solution and its configuration needs to be controlled by supramolecular complex formation with enantiopure counterions [3]. Here, we study the transfer of the underlying diastereomeric interaction to the photo-excited neutral quintet state, which couples to an accompanying ultrafast expansion of the metal-ligand distances.

I will conclude the talk by introducing a site-specific CD-label for time-resolved studies of ultrafast conformational dynamics in peptides and proteins [4]. In this way I hope to demonstrate how the measurement of broadband excited state CD spectra can open novel avenues in chiroptical spectroscopy and structural dynamics research.

[1] J. Meyer-Ilse et al., *Laser Photon. Rev.* **7**, 495 (2013)

[2] M. Oppermann et al., *Optica* **6**, 56 (2019)

[3] J. Lacour et al., *Angew. Chem. Int. Ed.* **37**, 2379 (1998)

[4] M. Oppermann & J. Spekowius et al., *J. Phys. Chem. Lett.* **10**, 2700 (2019)

Second Harmonic imaging of collagen 3D architecture in biological tissues: fast linear polarization resolution and Circular Dichroism

Margaux Schmeltz¹, Guillaume Ducourthial¹, Claire Teulon¹, Gaël Latour^{1,2}, Jean-Sébastien Affagard³, Jean-Marc Allain^{3,4}, Emmanuel Beaurepaire¹, Marie-Claire Schanne-Klein^{1*}

¹ LOB, Ecole Polytechnique, CNRS, INSERM, IPParis, 91128 Palaiseau, France

² Université Paris-Sud, CNRS, Université Paris-Saclay, F-91405 Orsay, France

³ LMS, Ecole Polytechnique, CNRS, IPParis, 91128 Palaiseau, France

⁴ Inria, Université Paris-Saclay, 91128 Palaiseau, France

e-mail : marie-claire.schanne-klein@polytechnique.edu

Collagen is the main component of connective tissues such as arteries, skin, bones or cornea. It is mainly organized as fibrils of various sizes and 3D distributions depending on the tissue. A defective collagen 3D organization or synthesis can lead to pathologies or tissue malfunctions, including poor mechanical behavior. *In situ* imaging of these 3D structures is therefore a major biomedical concern.

The gold standard technique for *in situ* 3D visualization of collagen fibrils in intact tissues is second harmonic generation (SHG) microscopy. Polarization-resolved SHG microscopy provides further information about the hierarchical organization of collagen, from the molecular structure to the mean orientation and degree of alignment of collagen fibrils at micrometer scale. In this work, a fast polarization-resolved SHG microscope is implemented using line-to-line switching of polarization by an electro-optical modulator and working in epidetection geometry. It is fast enough to obtain reliable orientations in dynamic collagen samples, while slow enough to get a good SHG signal to noise ratio in thick collagenous tissues. We demonstrate that it enables accurate quantitation of dynamic collagen reorganization in murine skin dermis during stretching mechanical assays.

Moreover, Circular Dichroism SHG (CD-SHG) microscopy is implemented to take advantage of collagen chirality to improve 3D visualization. It measures the normalized difference in the SHG signal obtained upon excitation by left versus right circular polarizations. In this work, we identify two major artifacts that may occur in CD-SHG experiments and we demonstrate that thorough optimization and calibration of the experimental setup are required for CD-SHG imaging. Finally, we successfully record CD-SHG images in human cornea sections and confirm that this technique efficiently reveals collagen fibrils oriented out of the focal plane.

Laser induced ultrafast spin dynamics in rare-earth / transition metals alloys by element- and time-resolved X-ray Magnetic Circular Dichroism

Nicolas Bergeard

*Institut de Physique et Chimie des Matériaux de Strasbourg, UMR 7504,
F-67000 Strasbourg, France
e-mail : nicolas.bergeard@ipcms.unistra.fr*

Femtomagnetism is one of the hottest topic in modern magnetism ever since its discovery in 1996 [BEA96]. The numerous theoretical and experimental works in this field of research aimed at identifying the microscopic mechanisms that allow the transfer of the angular momentum away from the spin degree of freedom during the laser induced ultrafast demagnetization. In this context, my main scientific interest concerns the investigation of laser induced ultrafast spin dynamics in rare-earth (RE) / transition metals (TM) alloys. Indeed, these alloys offers unique opportunities to address simultaneously the laser induced quenching of magnetic order for “localized” 4f and for “itinerant” 3d spins. Furthermore, their magnetic properties, such as the magnetic anisotropy, their temperatures of magnetic order, their magnetization, are controllably tunable with the concentration and the layer thickness. It is then possible to monitor the modification of the induced ultrafast dynamics of TM3d and RE4f spins in RE-TM alloys upon the controlled variation of these magnetic properties in order to challenge the predictions of theoretical models [KOO10, BAT10].

The investigations of RE 4f and TM 3d spin dynamics in RE-TM alloys require element- and time-resolved experiments. In my presentation, I will give an overview of our recent results obtained by mean of time-resolved X-ray Magnetic Circular Dichroism (TR-XMCD) experiments performed at large scale facilities either on the femtoslicing beamline at synchrotron BESSYII [BER14, FER17, FER19] or on the TEMPO beamline at synchrotron SOLEIL [SIL17].

[BEA96] Beaupaire et al. Phys. Rev. Lett 76, 4250, (1996)

[KOO10] Koopmans et al. Nature Mater. 9, 259 (2010)

[BAT10] Battiato et al. Phys. Rev. Lett 105, 027203 (2010)

[BER14] Bergeard et al. Nature Communications 5, 3466 (2014)

[FER17] Ferté et al. Phys. Rev. B 96, 134303 (2017) and Ferté et al. Phys. Rev. B 96, 144427 (2017)

[FER19] Ferté et al. Jour. Magn. And Magn. Mat. 485, 320 (2019)

[SIL17] Silly et al. J. Synchrotron Rad. 24, 886 (2017)

Light beams with orbital angular momentum : Generation, characterization and applications

Thierry Ruchon¹, Céline Chappuis¹, Romain Généaux^{1,2}, Thierry Auguste¹

1. LIDYL, CEA, CNRS, Université Paris-Saclay, CEA Saclay, 91191 Gif-sur-Yvette, France
2. Department of Physics, University of California, Berkeley, California 94720, United States

Light beams may carry both a spin and an orbital angular momentum (OAM). While the former is associated to their polarization state, the latter stems from the geometrical properties of their wavefront. In their prototypical form, beams with OAM have “donuts-like” intensity profile and helicoidal wavefront, carrying integral multiples of \hbar as angular momenta. Since their “rediscovery” in the late 90’s, beams with OAM of visible wavelengths have found innumerable applications in quantum optics, microscopy or information transfer. A major recent development was the generation of such beams with much smaller wavelengths – in the extreme ultraviolet (XUV) - using synchrotron sources, free electron lasers as well as high harmonic sources (HHG). In this latter case, it creates ultrashort XUV sources of beams with OAM, suited for time-resolved applications at femtosecond and attosecond time scales.

In this communication, we will first introduce the concept of OAM of light, starting from very first principles. We will then provide a selective overview of characterization schemes and applications of such beams. As a final part, we will review recent results aiming at providing beams with OAM in the XUV domain [1-4]

References

- [1] C. Chappuis, D. Bresteau, T. Auguste, O. Gobert, and T. Ruchon. High harmonic generation in an active grating. *arXiv e-prints*, 2018.
- [2] D. Gauthier, P. Rebernik Ribic, G. Adhikary, A. Camper, C. Chappuis, R. Cucini, L. F. DiMauro, G. Dovillaire, F. Frassetto, R. Généaux, P. Miotti, L. Poletto, B. Ressel, C. Spezzani, M. Stupar, T. Ruchon, and G. De Ninno. Tunable orbital angular momentum in high-harmonic generation. *Nature Communications*, 8:14971, apr 2017.
- [3] R. Généaux, A. Camper, T. Auguste, O. Gobert, J. Caillat, R. Taïeb, and T. Ruchon. Synthesis and characterization of attosecond light vortices in the extreme ultraviolet. *Nature Communications*, 7:12583, Aug 2016.
- [4] Romain Généaux, Céline Chappuis, Thierry Auguste, Samuel Beaulieu, Timothy T. Gorman, Fabien Lepetit, Louis F. DiMauro, and Thierry Ruchon. Radial index of laguerre-gaussian modes in high-order-harmonic generation. *Phys. Rev. A*, 95:051801, May 2017.